

ITA CORYZA ABC Gel

Inactivated aluminium hydroxide suspension vaccine
for the immunization of chickens against infectious coryza

Composition

ITA CORYZA ABC Gel inactivated vaccine contains a combination of *Avibacterium paragallinarum*, serotypes A, B and C, homogenized with aluminium hydroxide adjuvant and merthiolate as a preservative.

Indication

ITA CORYZA ABC Gel inactivated vaccine is recommended for the vaccination of breeder and layer type chicken flocks, against infectious coryza caused by *Avibacterium paragallinarum*.

Dosage & Administration

ITA CORYZA ABC Gel inactivated vaccine should be administered to breeder and layer-type chickens between 8 and 12 weeks of age. The vaccine must be injected subcutaneously (under the skin of the neck) or by intramuscular route (breast or thigh muscle) at the dose of 0.5 ml per bird. The program should be completed 4 to 6 weeks later, using an inactivated oil-adjuvant vaccine against infectious coryza.

Warnings & precautions

Do not mix the vaccine with other vaccines or drugs. If the vaccine is accidentally injected to the operator, urgent medical attention is required. Shake well before and during use. Do not freeze.

Satisfactory protection can be achieved only in well-developed, healthy birds. Use sterile injection equipment. Keep the vaccine at room temperature (15-25°C) for 3 to 4 hours before use. Check the accuracy of the syringe to ensure that the correct dose is delivered. Once a bottle has been opened, its contents must be used up within 24 hours, otherwise it should be discarded.

Storage

Store between +2°C and +8°C away from light and in a dry place. Keep all medicines out of reach of children.

Packaging

1000 doses (500 ml).

Manufactured by

Laprovvet, France

Marketed by

Navana Pharmaceuticals Ltd.
Rupshi, Rupganj, Narayanganj, Bangladesh

