

ITA ND+IBD

Inactivated, oil emulsion vaccine for the immunization of chickens against Newcastle Disease and Infectious Bursal Disease

Composition

ITA ND+IBD inactivated vaccine contains a combination of LaSota strain of Newcastle Disease virus and GP strain of the Infectious Bursal Disease virus in inactivated form, homogenized with oil adjuvant and merthiolate as a preservative.

Indication

ITA ND+IBD inactivated vaccine is recommended for the vaccination of breeder and layer type chicken flocks, previously immunized against Newcastle Disease and Infectious Bursal Disease with attenuated live virus vaccines. Progeny of vaccinates will have specific maternal antibodies against ND and IBD.

Dosage & Administration

ITA ND+IBD inactivated vaccine should be administered to breeder and layer-type chickens 3 to 4 weeks before the onset of lay, between 15 and 20 weeks of age. The vaccine must be injected subcutaneously (under the skin of the neck) or by intramuscular route (breast or thigh muscle) at the dose of 0.5 ml per bird.

Warnings & precautions

Do not mix the vaccine with other vaccines or drugs. If the vaccine is accidentally injected to the operator, urgent medical attention is required. Shake well before and during use. Do not freeze.

Satisfactory protection can be achieved only in well-developed, healthy birds. Use sterile injection equipment. Keep the vaccine at room temperature (15-25°C) for 3 to 4 hours before use. Check the accuracy of the syringe to ensure that the correct dose is delivered. Once a bottle has been opened, its contents must be used up within 24 hours, otherwise it should be discarded.

Side effects

None

Storage

Store between +2°C and +8°C away from light and in a dry place. Keep all medicines out of reach of children.

Packaging

1000 doses (500 ml).

Manufactured by

Laprovét, France

Marketed by

Navana Pharmaceuticals Ltd.
Rupshi, Rupganj, Narayanganj, Bangladesh

