

WSP
Tylos
(Tylosin tartrate BP)

Only for the use of animals

Composition

Each gram contains Tylosin tartrate BP 200 mg.

Pharmacology

Tylosin (Tylos) is a macrolide antibiotic that inhibits protein synthesis by binding with 50s ribosomal subunit of bacteria.

Indications Poultry

Mycoplasmosis, Chronic Respiratory Disease (CRD), Infectious synovitis, Pneumonia, Necrotic Enteritis etc. Livestock: Diseases caused by Streptococci, Staphylococci, Corynebacteria, Pasteurella etc. Pigs: Dysentery, Pneumonia, exudative epidermitis, arthritis and erysipelas.

Dosage and administration

Poultry: 2.5 g/ litter drinking water or 3-5 g/kg feed for 3-5 days. Calves: 4 g/50 kg body weight twice daily for 7-14 days. Swine: 1.25 g/ litter of drinking water for 7 - 10 days.

Or, as directed by Registered Veterinary Physician.

Contraindication

Tylosin (Tylos) should not use in animals with known hypersensitivity to tylosin or other macrolides & animals with hepatic disorders.

Warning & Precaution

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to animals not under treatment or wildlife.

Side Effects

It is safe in recommended dose. Occasionally diarrhea, articularia, pruritus, erythema may occur.

Drug Interaction

With Medicine: Should not use with aminoglycoside antibiotic.

With Food & Others: Not known.

Withdrawal period

Meat-1 day, Egg-0 (Zero) day

Storage

Store at a temperature below 30°C protected from light & keep in cool and dry place. Keep all medicines out of reach of children

Supply

100 g sachet.

Manufactured by:

Navana Pharmaceuticals Ltd.

Rupshi, Rupganj, Narayanganj, Bangladesh.

 **NAVANA PHARMA**
JAHURUL ISLAM COMPANY
Veterinary Division 