

Liquid
ENRO-10
(Enrofloxacin BP)

Only for the use in animals

Composition

Each ml contains Enrofloxacin BP 100 mg

Indications

For the prevention & treatment of respiratory and digestive tract infections caused by Mycoplasma, E. coli, Pasteurella, Salmonella etc. such as Mycoplasmosis, colibacillosis, Fowl Cholera, Salmonellosis etc. It is also indicated for the prevention & treatment of secondary bacterial infections followed by viral infections such as Ranikhet, Gumboro etc.

Dosage & Administration

Prevention: 1 ml/4-8 L drinking water for 3-5 days.

Treatment: 1 ml/2 L drinking water for 3-5 days. Or as directed by the registered Veterinary Physician.

Contraindication

Contraindicated in birds hypersensitive to enrofloxacin.

Drug Interaction

Should not use enrofloxacin along with chloramphenicol, macrolides & tetracyclines.

Withdrawal Period

Meat: 3 days Egg: Should not use in bird producing eggs for human consumption.

Storage

Store at a temperature below 30°C protected from light & keep in cool and dry place. Keep all medicines out of reach of children

Packing

100 ml & 1L bottle

Manufactured by:

Navana Pharmaceuticals Ltd.

Rupshi, Rupganj, Narayanganj, Bangladesh.

