

**ITA - NEW (ND)**  
Inactivated, oil emulsion vaccine  
for the immunization of chickens against Newcastle Disease

**Composition**

ITA - NEW (ND) inactivated vaccine contains LaSota strain of Newcastle Disease virus in inactivated form, homogenized with oil adjuvant and merthiolate as a preservative.

**Indication**

ITA - NEW (ND) inactivated vaccine is recommended for the vaccination of breeder and laying-type chicken flocks, previously immunized against Newcastle Disease (ND) with an attenuated live virus vaccine.

**Dosage & Administration**

ITA - NEW (ND) inactivated vaccine should be administered to breeder and laying-type chickens 3 to 4 weeks before the onset of lay, between 15 and 20 weeks of age. The vaccine must be injected subcutaneously (under the skin of the neck) or by intramuscular route (breast or thigh muscle) at the dose of 0.5 ml per bird.

**Warnings & precautions**

Do not mix the vaccine with other vaccines or drugs. If the vaccine is accidentally injected to the operator, urgent medical attention is required. Shake well before and during use. Do not freeze.

Satisfactory protection can be achieved only in well-developed, healthy birds. Use sterile injection equipment. Keep the vaccine at room temperature (15-25°C) for 3 to 4 hours before use. Check the accuracy of the syringe to ensure that the correct dose is delivered. Once a bottle has been opened, its contents must be used up within 24 hours, otherwise it should be discarded.

**Side effects**

None

**Storage**

Store between +2°C and +8°C away from light and in a dry place. Keep all medicines out of reach of children.

**Packaging**

1000 doses (500 ml).

**Manufactured by**

Laprovét, France

**Marketed by**

Navana Pharmaceuticals Ltd.  
Rupshi, Rupganj, Narayanganj, Bangladesh

