

**Solution**  
**ACTIVATE WD MAX**

Only for the use of animals

**Composition**

Total acid (min) Formic acid 34%, Propionic acid 20% and HMTBa (2-Hydroxy-4-Methylthio Butanoic acid) 35%.

**Description**

ACTIVATE WD MAX is used in drinking water for the control and destruction of environmental pathogens (*Salmonella*, *E.coli*, *Campylobacter*, etc...). ACTIVATE WD MAX has a nutritional effect due to the supply of methionine, which is known to aid in the detoxification and stimulation of the hepatic system.

**Indication**

Activate WD MAX is composed of Gut environment modifiers designed to improve Feed and Water quality. It prevents the colonization of the proximal small intestine by harmful bacteria avoiding bacterial development in water and feed. Acidification of the feed and drinking water by Activate WD MAX reduces its microbial load. The use of Activate WD MAX is a preventive measure to control *Salmonella*, *E.coli*, *Campylobacter* etc.

**Dosage & Administration**

*Poultry:* 1ml / 2-4 L drinking water for 5 days.

Or as directed by the registered veterinary physician or nutritionist.

*Aqua:* For 3ft depth:

-pH above 9: 10ml/Decimal

-pH 8.5-9: 6 ml/Decimal

**Warnings & precautions**

please use gloves while handling.

**Side effects**

Common: No side effects observed in recommended dose.

Rare: None

**Withdrawal periods**

Nil

**Storage**

Store below 30°C away from light and in a dry place. Keep all medicines out of reach of children.

**Packing**

198 L drum

**Manufactured by**

Novus International Co. Ltd.

Germany

**Marketed by**

Navana Pharmaceuticals Ltd.

Rupshi, Rupganj, Narayanganj, Bangladesh

