

Injection

# Phenira **Vet**

Pheniramine Maleate BP

**Composition:**

Each ml contains Pheniramine Maleate BP 22.75 mg.

**Pharmacology**

Pheniramine Maleate is the maleate salt form of pheniramine, an alkylamine derivative with antihistaminic and vasodilatory properties. Pheniramine maleate binds to histamine H1 receptors, thereby inhibiting phospholipase A2 and production of endothelium-derived relaxing factor, nitric oxide.

**Indications:**

Itching of unknown genesis and of various localizations, eczema, dermatitis, urticaria, skin oedema, insect bites, photodermatitis, rhinitis, tail eczema in horses, stomatitis, toxic hoof corns and inflammation of the hooves of cattle, serum shock, paresis during pregnancy, puerperal toxemia and secondary retention, pulmonary oedema in cattle, pulmonary emphysema in horses etc. The efficacy of **Phenira Vet** has also been demonstrated in drug allergies, anaphylactic shock, laminitis and bloat in ruminants due to histamine atony of the rumen and also in the treatment of acute septic metritis in cows in conjunction with antibacterial therapy.

**Dosage and Administration:**

Cattle, Buffalo & Horse: 5 - 10 ml per day.

Goat & Sheep: 0.5 - 1 ml per day.

Phenira Vet may be administered intramuscularly or intravenously. The onset of action depends on the mode of administration. Mostly there is an immediate effect on intravenous administration; on intramuscular administration, it takes about 10-20 minutes for the effect to set on.

**Or as directed by the registered Veterinary Physician.**

**Contraindication:**

The drug is contraindicated in animal hypersensitive to active ingredient.

**Warnings & precautions**

The dosage should be adjusted individually. If possible small doses are given first because it has often been shown that anti-allergic drugs are effective in comparatively small dosage.

**Side effects**

Common: In therapeutic doses, slight insomnia, drowsiness may develop & may feel fatigue. Injection may be irritant and transient hypotension or stimulation of the CNS.

Rare: Not Known.

**Use in Pregnancy & Lactation:**

Safety for the use of Phenira Vet injection during pregnancy has not been established. This product should only be used when clearly needed and potential benefit outweighs the potential unknown risks to the fetus. Use during lactation is not recommended.

**Overdose**

The drug can be administered 2-3 times daily, depending on the severity of the disease.

**Withdrawal Period:**

Meat- 7 days & Milk- 2 days

**Storage**

Store in a cool, dry place & keep away from light.

Keep out of the reach of children.

**Packing**

10ml Vial, 100ml Vial.

**Manufactured by**

Navana Pharmaceuticals Ltd.

Rupshi, Rupganj, Narayanganj, Bangladesh

